




**No. 3** March 2009 **EUROPE**

**CVBD<sup>®</sup>**

**DIGEST**

www.cvbd.org



# Climate change – Is it paving the way for the expansion of CVBD?

Cutting-edge information  
brought to you by  
the CVBD World Forum



## Introduction

When Bayer Animal Health called for the 1<sup>st</sup> International CVBD Symposium in 2006, it was the first and initial step to addressing the global threat of canine vector-borne diseases (CVBD). This was based on the belief that canine vector-borne diseases should be treated as one topic and dealt with on a global level in an interdisciplinary way. During the past years, CVBD have become a global issue and even sparked public interest. Many of the parasite-transmitted diseases affect humans as well as animals. As man's best friend, dogs play an important role as they are affected to a high extent by zoonotic pathogens, even serving as a host for some of them.

At the first symposium, the participants agreed to form the CVBD World Forum. Besides gathering knowledge, the main task of this group of international experts has been to raise awareness for the specific regional risks of CVBD and to foster preventative measures. For this reason, the CVBD World Forum created a website ([www.cvbd.org](http://www.cvbd.org)) to provide the veterinary practitioners with cutting-edge and clinically relevant scientific information on CVBD.

In the CVBD Digest, relevant findings from CVBD symposia are presented periodically to veterinary practitioners around the globe. During the three symposia so far, the impact of climate change on CVBD has been an important issue. While global warming is more or less accepted as a fact by both experts and the public, the effect on parasites and their transmitted diseases is still the subject of heavy debate. There is also a local component of climate change that strongly influences the regional situation of CVBD. Moreover, veterinarians should consider other man-made factors such as pet travel, mobility or import – all of which could raise the CVBD threat for dogs in many, if not most parts of the world.

Cutting-edge information brought to you by the CVBD World Forum

## Climate change – Does it pave the way for an expansion of CVBD?

**Author: Friederike Krämer**

Institute for Parasitology, University of Veterinary Medicine  
Hanover, Germany


**Parasite-transmitted diseases are one of the major contributors to the global burden of disease in humans and animals. Climate change is often held responsible for the spread of parasitic vectors such as ticks, fleas, mosquitoes and sand flies, and their transmitted pathogens – leading, in the case of the dog, to so-called canine vector-borne diseases (CVBD). Currently, there is insufficient data available to prove if climate change is a major driving force for vector and disease expansion. Other reasons, such as ecological, demographic and socio-economic factors e.g. pet travel into and pet import from endemic areas, also play a role in this development. Whatever the reasons might be for possible disease expansion, prevention has to be focused on educating dog owners who are responsible for individual parasite protection as well as for risk-averse behaviour, e.g. regarding pet travel. Broad-spectrum vector control should be practiced by using parasiticides that repel and kill blood feeders in order to minimize the risk of CVBD-pathogen transmission.**

### Climate change and vector-borne diseases

Climate change is defined as a statistically significant variation in either the mean state of the climate or in its variability, persisting for an extended period of time. Meanwhile there is good evidence that climate is changing. The global average land and sea surface temperature has increased by  $0.6 \pm 0.2$  °C

since the mid-19<sup>th</sup> century.<sup>1</sup> Furthermore, the Intergovernmental Panel of Climate Change (IPCC) 4<sup>th</sup> Assessment Report states that “most of the observed increase in global average temperatures since the mid-20<sup>th</sup> century is very likely due to the observed increase in anthropogenic greenhouse gas concentrations, and that it is likely that there has been significant anthropogenic warming over the past 50 years averaged over each continent (except Antarctica)”.<sup>2</sup> For the future, this process might even be accelerated (Fig1). However, besides global climate changes, regional and local climate development as well as microclimate changes have to be considered, especially since they are also of major importance for vector-borne disease development (see below).

An increasing number of studies and publications try to determine whether, and how, climate change is affecting vector-borne diseases. Most intermediate hosts and vectors and the majority of reservoirs have well defined geographic distributions resulting from evolutionary processes that condition the disease they transmit or shelter. Within its area of distribution, each species finds more or less suitable conditions for its development and multiplication and thus for disease transmission. Any events that modify the climate or environment affect the epidemiology of a disease.<sup>3</sup> Global climate change might expand the distribution of vector-borne pathogens in both time and space, thereby exposing host populations to longer transmission seasons, and immunologically naïve populations to newly introduced pathogens.<sup>4</sup> However, caution is necessary when interpreting these predictions.



With respect to disease transmission, changes in the following properties are considered to be most important: (i) survival and reproduction rates of vectors, in turn determining their distribution and abundance; (ii) intensity and temporal pattern of vector activity (particularly biting rates) throughout the year; and (iii) rates of development, survival and reproduction of pathogens within vectors.<sup>5</sup> Additionally, the ability of disease-transmitting vectors to react to climate changes via active or passive dispersal also has to be considered (Tab. 1).

### The role of temperature

The surrounding temperature directly influences the physiology of the cold-blooded vectors and the pathogens they submit. In many cases the pathogen only develops within the vector (outer incubation) within a certain temperature range, even though both may tolerate more extreme temperatures. For example, development of the so-called heartworm *Dirofilaria immitis* in its mosquito host needs an average temperature of at least 18 °C over a 30-day

## INFO BOX 1

### HEARTWORM DISEASE IN THE UK?

Studies have shown that summer temperatures in certain parts of the UK and during particular years can facilitate the incubation of *Dirofilaria immitis* larvae in the mosquito intermediate host<sup>25</sup>, with transmission being dependent on the presence of sufficient numbers of susceptible mosquitoes and a large enough population of infected dogs. It is suggested in turn that summer conditions as described above may occur in three out of five summers by 2080 in the UK.<sup>26</sup> Similar facilitating temperature conditions for at least one heartworm generation have also been detected in the Czech Republic.<sup>27</sup>

Vector	Major disease	Active dispersal	Passive dispersal
<i>Ixodes</i> ticks	Lyme borreliosis, tick-borne encephalitis	very limited powers of active dispersal	common and wide ranging, due to long periods of feeding: attachment to birds and large mammals
Mosquitoes	malaria, (diro)filariasis, dengue fever, yellow fever, West Nile fever (WNV)	able to disperse and reproduce relatively rapidly to exploit newly suitable areas	common and wide ranging: often transported by human transportation, including inter-continental air crafts and cargo
Sand flies	leishmaniosis	limited flight ranges	highly unlikely
Fleas	bartonellosis	limited dispersal, inbetween couple of centimeters as emerging adults	common and wide ranging, due to infestation of mammals

Tab. 1: Potential vector responses to climate change (modified after reference 5)

	Mosquito	Pathogen
transmission increased due to	abbreviated developmental time of larvae and pupae abbreviated gonotrophic cycle more frequent oviposition increased biting frequency	faster replication  abbreviated outer incubation time (pathogen within the vector)
transmission reduced due to	abbreviated life time	

Tab. 2: Effects of temperature increase on vector (i.e. mosquito), pathogen and frequency of transmission<sup>24</sup>

period (see also [Info box 1](#)). Below 18 °C larval development is retarded, but partial development still occurs. The lower threshold temperature, below which larval development does not take place, is generally agreed to be 14 °C for *D. immitis*.<sup>6,7,8,9</sup>

For the protozoan *Leishmania* sp. a minimum of 10 °C is necessary to initiate the invertebrate cycle, and the rising temperature is continuously shortening the time needed for completion, which is at least 6 days for *Leishmania infantum*.<sup>10,11</sup> The 10 °C year isotherm is furthermore limiting the occurrence of the *Leishmania* vector *Phlebotomus* sp. During recent years, *Phlebotomus* sand flies have been known to expand e.g. to northern regions in Italy<sup>12</sup> (see also [Info box 2](#)). Thus, both vector and pathogen might be influenced in their geographical distribution by rising temperatures.

## INFO BOX 2 EXPANSION OF SAND FLIES TO NORTHERN ITALY


Compared to historical data, *Phlebotomus perniciosus* and *P. neglectus* have increased in density and expanded their geographic range in Northern Italy. Northern continental Italy is now focally endemic for visceral leishmaniosis (VL) and a moderate risk of human disease does exist, although the intensity of transmission seems to be lower than in traditional settings of Mediterranean VL.<sup>12</sup>

In general, higher temperatures usually abbreviate the development time of the outer incubation of the pathogen, but also abbreviate the survival time of the vector with oppositional effect on the frequency of transmission.<sup>13</sup> The effects of the temperature rise are considered to cause an increase of the pathogen transmission rate in the first place (Tab. 2). For ticks both the delay period and mortality vary geographically and seasonally with climate. While tick birth and development rates increase with temperature, tick mortality rates increase with moisture stress.<sup>14</sup>

### Microclimate affecting ticks

Climate change might not only have a distinct effect on the development cycle of a vector. Through changing precipitation and even seasons, climate change also alters the microclimate as well as tick biology and feeding behaviour.

*Ixodes* ticks, for example, are very susceptible to desiccation while questing for hosts in the vegetation. For passive and active atmospheric water uptake, they periodically return to moist conditions at the base of the vegetation to restore their fluid content.<sup>15,16,17,18</sup> It has been observed that under increasingly dry conditions the number of unfed nymphal *Ixodes ricinus* questing in upper layers of the herbage decreased, whereas the rate of fat use and the number of nymphs feeding on small rodents both increased. For *Ixodes* larvae, very few larvae quested or fed on rodents in dry conditions, but many more did so once the humidity increased, suggesting that larvae escape desiccation by becoming quiescent. The ratio of larvae to nymphs feeding on rodents thus increased with rising humidity, contributing to the seasonal and geographical variation in disease transmission dynamics.<sup>19</sup> Nevertheless, the authors admit that the behaviour described above may be different in ticks from different regions which might be adapted and accli-





## INFO BOX 3

### NORTHWARD SPREAD OF *IXODES* TICKS IN SWEDEN

Results of a questionnaire in Sweden show a greater occurrence of *Ixodes* ticks between the early 1980s and early 1990s, indicating a northward shift in the latitudinal limit for reported tick occurrence.<sup>25</sup> Furthermore, the incidence of human TBE in Sweden approximately doubled for the period 1984–1994, compared with 1960–1983. A relatively mild climate of the 1990s in Sweden was suggested as one of the probable

primary reasons for the observed increase of density and geographic range of the Castor Bean tick *I. ricinus*.<sup>26</sup> An increase in the roe deer population from the mid-1980s likely also influenced the tick density.<sup>26</sup> It has to be further judged whether the better recollection of ticks and the possibility of increased awareness, detection and reporting of cases also have to be accounted for in this increase.<sup>5</sup>

matized to their particular microclimate. Despite this possible influence of climate on ticks, changing climatic factors that explain both the spatial variation and the temporal patterns in tick-borne disease 'emergence' have not yet been identified in detail. Retrospective climate evaluation in the UK e.g. did not reveal obvious differences between sites where human tick-borne encephalitis (TBE), as one of the best studied tick-borne diseases, did or did not 'emerge'.<sup>14</sup> But field studies by Ogden and colleagues<sup>20</sup> in Canada at least supported temperature and tick immigration based risk maps (Fig. 2) for the mere northward expansion of *Ixodes scapularis* (see also Info box 3).

#### Impact of other factors

The ecology and epidemiology of vector-borne diseases are affected by the interrelations between four major factors comprising the pathogen, the host (human or animal), the vector and the environment. Besides atmospheric and climatic changes, alternative 'background' socio-economic, demographic and ecological effects (see also Info box 4 and 5) are plausible enough to cast some doubt on the role of climate change<sup>21</sup>.

Some of these effects facilitating a spread of vectors and pathogens are illustrated in more detail below:

- (i) extended outdoor activities leading to higher exposures, either by an increase in leisure time (increased living standard) or by an increase in outdoor work and use of natural resources (decreased living standard);

## INFO BOX 4

### IMPORTANT DRIVERS FOR VECTOR-BORNE PARASITE EMERGENCE AND SPREAD<sup>27</sup>

- habitat changes
- alterations in water storage and irrigation habits
- pollution
- development of insecticide and drug resistance (e.g. *Anopheles* sp. and DDT)
- globalization
- increase in international trade, tourism and travel
- immunodeficient hosts (e.g. HIV infection in humans)
- war and civil unrest
- global and governmental management failure

- (ii) altered farming conditions, increased scrubland, urbanization with poor hygienic measures;
- (iii) increased (pet) travel, increased pet import into non-endemic regions

Concerning (i) and (ii), a well-studied example is the increase in tick-borne diseases (representative here tick-borne encephalitis [TBE]) in man in the Baltic States at the end of the communist rule in Eastern Europe<sup>28</sup> (see also [Info box 5](#)).

In the case of pet travel and the import of dogs from endemic areas (iii), 20,000 imported *Leishmania*-infected dogs are estimated to be living in Germany. These could act as a reservoir if the appropriate *Phlebotomus* species became endemic in this country or if the autochthonous species *P. mascitii* is a capable vector (see also [Info box 6](#)).

## INFO BOX 5

### INCREASE OF HUMAN TBE IN THE BALTIC STATES

Randolph<sup>28</sup> identified the following enhancing conditions for an increase of tick-borne encephalitis (TBE) incidence at the end of the communist rule in Eastern Europe: a decline of agriculture with changes in land use and cover, causing large areas of abandoned land that are increasingly for rodent transmission hosts; increased unemployment and poverty with subsequent in-

crease of subsistence farming; increased material welfare in other parts of the population with regular weekend trips to country dachas; a decline in national cattle herds; reduced maintenance of grassed fields and increased goat numbers, offering better feeding opportunities to ticks in marginal lands and increasing the risk of alimentary TBE transmission route.

## INFO BOX 6

### IMPORTED LEISHMANIOSIS IN CENTRAL EUROPE

The testing of 291 dogs imported from or returning to Germany, Switzerland or Austria from Southern Europe (Spain, Italy, Greece, Turkey, France, Malta, Portugal and others) between February and October 2003 resulted in 111 dogs (38%) classified serologically positive for *Leishmania*.<sup>29</sup> Of these 111 dogs, 103 had been imported and eight were travelling dogs, showing the high importance of dog importing

for disease entry. Nevertheless, surveys from the Netherlands estimate that some 58,000 travelling Dutch dogs are at risk of exposure to a *Leishmania* infection in Southern Europe each year<sup>30</sup>, based on a total population of 1,200,000 dogs in the country. An increase of endemic areas was e.g. recorded in South-western France between 1987 and 2004<sup>31</sup>, thus setting the travelling dog at a higher risk.



## Conclusion

Undoubtedly, changing climatic conditions and their alterations in temperature and humidity do have an influence on disease-transmitting vectors and their surrounding microclimate. However, the availability of hosts is a major limiting factor for the spread of vector and pathogen.

Although biological reasons seemingly support the theory that anticipated climate change will cause a spread of vector-borne diseases, ecological and socioeconomic factors have in the past proven to be stronger driving forces for the spread of infectious diseases than climate.<sup>13</sup> At a continental scale, climate may be a good predictor of disease distribution (e.g. Lyme disease in the USA<sup>22</sup>), but on the 'local scale' other environmental factors are likely to be more important.<sup>5</sup> To prove the individual effect of each factor, more data, new approaches and intensive studying are necessary. A step into that direction is disease distribution mapping with the help of remote sensing and GIS technology. Long term data over decades in extended areas are needed. The often missing strong evidence of

climate change on vector-borne diseases has to be seen as an 'absence of evidence', rather than an 'evidence of absence', as stated by Kovats and colleagues.<sup>5</sup>

Transmission of clinically important canine vector-borne organisms to new regions, either due to climatic factors or due to dog travel and import, is of particular concern to veterinary clinicians, as the diseases caused by these organisms might be unfamiliar to them and may go undiagnosed.<sup>23</sup> Therefore, when it comes to disease prevention, education of the dog owner regarding individual parasite protection and risk-adverse behaviour is still the method of choice. Broad-spectrum ectoparasiticides on the basis e.g. of a monthly spot-on application can protect against ticks, fleas, sand flies, mosquitoes and stable flies. If these remedies also provide a repellent effect, they will further reduce the risk of CVBD-pathogen transmission via the aforementioned vectors which can cause dangerous diseases in dogs. Such preventative measures should be frequently applied in endemic areas to control a disease as well as in dogs which are brought there from non-endemic regions.

## References

1. Intergovernmental Panel on Climate Change (IPCC) (2001): Climate change 2001: The scientific basis. Summary for policy-makers. Cambridge University Press, Cambridge, UK, 1–20
2. Intergovernmental Panel on Climate Change (IPCC) (2007): Climate change 2007: Synthesis report. Summary for policy-makers. Cambridge University Press, Cambridge, UK, 1–22
3. Mouchet, J., Carnevale, P. (1997): [Impact of changes in the environment on vector-transmitted diseases.] *Sante* 7, 263–269
4. Patz, J.A., Reisen, W.K. (2001): Immunology, climate change and vector-borne diseases. *Trends Immunol.* 22, 171–172
5. Kovats, R.S., Campbell-Lendrum, D.H., McMichael, A.J., Woodward, A., Cox, J.S. (2001): Early effects of climate change: do they include changes in vector-borne disease? *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 356, 1057–1068
6. Genchi, C., Rinaldi, L., Cascone, C., Mortarino, M., Cringoli, G. (2005): Is heartworm disease really spreading in Europe? *Vet. Parasitol.* 133, 137–148
7. Knight, D.H., Lok, J.B. (1998): Seasonality of heartworm infection and implications for chemoprophylaxis. *Clin. Tech. Small Anim. Pract.* 13, 77–82
8. Lok, J.B., Knight, D.H. (1998): Laboratory verification of a seasonal heartworm model. In: Seward, R.L.: Recent Advances in Heartworm Disease. Symposium '98, Am. Heartworm Soc., Batavia, USA, 15–20
9. Slocombe, J.O.D., Surgeoner, G.A., Srivastava, B. (1989): Determination of heartworm transmission period and its use in diagnosis and control. *Proc. Heartworm Symposium 1989*, Am. Heartworm Soc., Washington D.C., USA, 19–26
10. Molyneux, D.H., Killick-Kendrick, R. (1987): Morphology, ultrastructure and life cycles. In: Peters, W., Killick-Kendrick, R. [eds.]: *The Leishmaniasis in Biology and Medicine*. Academic Press, London, UK, Vol. 1, 121–176

11. Killick-Kendrick, R. (1990): The life cycle of *Leishmania* in the sandfly with special reference to the form infective to the vertebrate host. *Ann. Parasitol. Hum. Comp.* 65 (Suppl. 1), 3742
12. Maroli, M., Rossi, L., Baldelli, R., Capelli, G., Ferroglia, E., Genchi, C., Gramiccia, M., Mortarino, M., Pietrobelli, M., Gradoni, L. (2008): The northward spread of leishmaniasis in Italy: evidence from retrospective and ongoing studies on the canine reservoir and phlebotomine vectors. *Trop. Med. Int. Health* 13, 256–264
13. Ebert, B., Fleischer, B. (2005): [Global warming and spread of infectious diseases.] *Bundesgesundheitsbl. Gesundheitsforsch. Gesundheitssch.* 48, 55–62
14. Randolph, S.E. (2004): Evidence that climate change has caused 'emergence' of tick-borne diseases in Europe? *Int. J. Med. Microbiol.* 293 (Suppl. 37), 5–15
15. Lees, A.D., Milne, A. (1951): The seasonal and diurnal activities of individual sheep ticks (*Ixodes ricinus* L.). *Parasitol.* 41, 189–208
16. Knülle, W., Rudolph, D. (1982): Humidity relationships and water balance of ticks. In: Obenchain, F.D., Galun, R. [eds.]: *Physiology of Ticks*. Pergamon Press, Oxford, UK, 43–70
17. Needham, G.R., Teel, P.D. (1991): Off-host physiological ecology of ixodid ticks. *Ann. Rev. Entomol.* 36, 659–681
18. Sonenshine, D.E. (1991): *Biology of Ticks*. Oxford University Press, Oxford, UK, Vol. 1
19. Randolph, S.E., Storey, K. (1999): Impact of microclimate on immature tick-rodent host interactions (Acari: Ixodidae): implications for parasite transmission. *J. Med. Entomol.* 36, 741–748
20. Ogden, N.H., St-Onge, L., Barker, I.K., Brazeau, S., Bigras-Poulin, M., Charron, D.F., Francis, C.M., Heagy, A., Lindsay, L.R., Maarouf, A., Michel, P., Milord, F., O'Callaghan, C.J., Trudel, L., Thompson, R.A. (2008): Risk maps for range expansion of the Lyme disease vector, *Ixodes scapularis*, in Canada now and with climate change. *Int. J. Health Geogr.* 7, 24, doi: 10.1186/1476-072X-7-24
21. Korenberg, E.I., Kovalevskii, Y.V. (1999): Main features of tick-borne encephalitis eco-epidemiology in Russia. *Zentralbl. Bakteriol.* 289, 525–539
22. Glass, G.E., Schwartz, B.S., Morgan, J.M. III, Johnson, D.T., Noy, P.M., Israel, E. (1995): Environmental risk factors for Lyme disease identified with geographic information systems. *Am. J. Public Health* 85, 944–948
23. Irwin, P.J. (2002): Companion animal parasitology: a clinical perspective. *Int. J. Parasitol.* 32, 581–593
24. Mellor, P.S., Leake, C.J. (2000): Climatic and geographic influences on arboviral infections and vectors. *Rev. Sci. Tech.* 19, 41–54
25. Tälleklint, L., Jaenson, T.G. (1998): Increasing geographical distribution and density of *Ixodes ricinus* (Acari: Ixodidae) in central and northern Sweden. *J. Med. Entomol.* 35, 521–526
26. Lindgren, E., Tälleklint, L., Polfeldt, T. (2000): Impact of climatic change on the northern latitude limit and population density of the disease-transmitting European tick *Ixodes ricinus*. *Environ. Health Perspect.* 108, 119–123
27. Harrus, S., Baneth, G. (2005): Drivers for the emergence and re-emergence of vector-borne protozoal and bacterial diseases. *Int. J. Parasitol.* 35, 1309–1318
28. Randolph, S.E. (2008): Tick-borne encephalitis incidence in Central and Eastern Europe: consequences of political transition. *Microbes Infect.* 10, 209–216
29. Mettler, M., Grimm, F., Naucke, T.J., Maasjost, C., Deplazes, P. (2005): [Canine leishmaniosis in Central Europe: retrospective survey and serological study of imported and travelling dogs.] *Berl. Münch. Tierärztl. Wochenschr.* 118, 37–44
30. Teske, E., van Knapen, F., Beijer, E.G., Slappendel, R.J. (2002): Risk of infection with *Leishmania* spp. in the canine population in the Netherlands. *Acta Vet. Scand.* 43, 195–201
31. Bourdeau, P. (2008): Canine vector-borne diseases in France: information obtained from veterinary clinics in national surveys. *Proc. 3rd Canine Vector-Borne Disease (CVBD) Symposium, Wiesbaden, Germany, April 16.–19., 2008*, 78–84
32. Greer, A., Ng, V., Fisman, D. (2008): Climate change and infectious diseases in North America: the road ahead. *CMAJ* 178, 715–722

\* Member of the CVBD World Forum





