


## VACCINATION IN DRINKING WATER

### Step-by-step administration guide



## Reconstitution of the oral vaccine



**For 100 piglets, use 2 vials and fill up with mineral water to the 200 ml mark**

**Elanco**

1. L. Bélanger *et al.* 2017. *In vitro* investigation on the compatibility of Coliprotec® F4/F18, live oral *E. coli* vaccine against post-weaning diarrhoea, and Avibblue® water stabilizer. Proceedings of the 9th European Symposium of Porcine Health Management, 3-5 May Prague, Czech Republic, p 316.

Coliprotec® F4/F18 lyophilisate for oral suspension for pigs. Each dose of vaccine contains  $1.3 \times 10^8$  to  $9.0 \times 10^8$  CFU of live non-pathogenic *Escherichia coli* O8:K87 (F4ac) and  $2.8 \times 10^8$  to  $3.0 \times 10^9$  CFU of live non-pathogenic *Escherichia coli* O141:K94 (F18ac). Legal category POM-V (UK). For further information call Elanco Animal Health on +44(0)1256 353131 or write to: Elanco Animal Health, Lilly House, Priestley Road, Basingstoke, Hampshire RG24 9NL, or consult the product SPC. Advice should be sought from the Medicine Prescriber. **Use medicines responsibly** [www.noah.co.uk/responsible](http://www.noah.co.uk/responsible). Coliprotec is a registered trademark of Prevet Microbia Inc, used under license by Elanco. Elanco and the diagonal bar logo are trademarks of Elanco or its affiliates. © 2020 Elanco or its affiliates. PM-UK-20-0183.

# Administration of the vaccine before weaning



TROUGH / BOWL

4

In a 10 L bucket:

1. Add 5 g or 1 teaspoon of *AviBlue*®1
2. Add 10 L of water




5

- 1 Transfer 1 L of solution



- 2 Add the number of doses corresponding to the number of piglets in the litter (a single 2 ml dose per piglet)



Transfer 1 L solution with the vaccine into the water bowl (should be clean and free from food)

# Administration of the vaccine after weaning


4

In a 10 L bucket:


1. Add 5 g or 1 teaspoon of *AviBlue*®1
2. Add 10 L of water

1 bucket of 10 L = 60 piglets at 21 days of age


1 bucket of 10 L = 40 piglets at 28 days of age



5



- 2 Add the corresponding number of doses for number of piglets to be vaccinated (1 single dose of 2 ml per piglet)



- 3 Distribute in the water bowl/trough

	Weaning 21 days	Weaning 28 days
Piglets/ box	Water/box (in Litres)	
15	2.55	3.5
20	3.5	4.6
25	4.75	5.75
30	5	7