

WARNING

KEEP OUT OF REACH OF CHILDREN

Dangerous to the Environment

Corrosive

PLEASE READ THE LABEL COMPLETELY BEFORE USE

Hazard Classification: 6.3A, 6.9B, 8.3A, 9.1C

Safety Directions: May cause skin irritation. Avoid skin contact. May cause eye damage. Avoid contact with eyes. Wear eye and face protection. Cyromazine possibly may affect blood and the haemopoietic system. Acetic acid possibly may damage lungs following inhalation. Handle with care.

First Aid: If skin or hair contact occurs remove contaminated clothing and flush skin and hair with running water. If eye contact occurs hold eyes open and rinse cautiously with water for several minutes. Contact the National Poisons Centre **0800 POISON (0800 764766)** or a doctor immediately.

Environmental Protection: Harmful to aquatic life. Avoid contamination of any water supply with product or empty container.

Emergency Response: In case of spillage, wear appropriate protective clothing and prevent material from entering waterways. Absorb spills with inert material and place in waste containers. Wash the area with water and absorb with further inert material. Dispose of waste safely.

Disposal: Preferably dispose of the product by use. Otherwise dispose of product and packaging at an approved landfill or other approved facility. Triple rinse container with water and dispose of rinsate away from waterways. Crush or puncture and bury in a suitable landfill. Do NOT use container for any other purpose.

*For further details see the Safety Data Sheet

VETRAZIN™

SPRAY-ON

DIRECTIONS FOR USE


Apply VETRAZIN SPRAY-ON through a suitable clean applicator gun. The product should be sprayed on in bands with the gun held 20 cm above the wool during application. VETRAZIN SPRAY-ON should be applied onto those areas where maximum protection against flystrike is required. VETRAZIN SPRAY-ON applied between the rump and shoulders should be sprayed on in even bands as shown (see diagram). The product sprayed onto the rump should be applied in an arc over the area shown (see diagram). The dose rate and number of bands to be applied are shown in the table.

Note: The dose volume applied should be based on the weight of the heaviest sheep in the mob.

Application and dose rate of VETRAZIN SPRAY-ON for the protection of all the commonly struck areas of the sheep.

Application and dose rate of VETRAZIN SPRAY-ON at docking for the prevention of flystrike.

Apply between 5 mL and 10 mL over the crutch ensuring that sufficient product to cover the area to be protected is applied (see diagram).



	2 bands Sheep between 11-20 kg		3 bands Sheep between 21-40 kg		4 bands Sheep between 41-60 kg		5 bands Sheep between 61-70 kg	
Liveweight (kg)	11-15	16-20	21-30	31-40	41-50	51-60	61-70	
Number of bands	2	2	3	3	4	4	5	
Volume per band (mL)	7.5	10	10	13.3	12.5	15	14	
Total volume applied (mL)	15	20	30	40	50	60	70	
Sheep treated per 25 L	1665	1250	832	625	500	415	355	

By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

It is an offence for users of this product to cause residues exceeding the relevant MRL in the New Zealand (Maximum Residues Limits of Agricultural Compounds) Food Standards.

WITHOLDING PERIODS

1. **MEAT:** Sheep producing meat or offal for human consumption must not be sold for slaughter either during treatment or within 7 days of cessation of the last treatment.

2. **MILK:** Milk intended for sale for human consumption must be discarded during treatment for not less than 35 days following the last treatment.

WOOL RESIDUES

Ensure the time between dipping and subsequent shearing is at least two months, except in the case of spot treatment of flystrike. For advice on your dipping programme contact your supplier.

ACTIVE INGREDIENT

VETRAZIN SPRAY-ON contains 60 g/litre cyromazine in the form of an aqueous concentrate.

GENERAL INFORMATION

VETRAZIN SPRAY-ON is an easy to use spray-on, which will give up to 6 weeks residual protection against flystrike caused by all the commonly found strike flies including the Australian blowfly (*Lucilia cuprina*). VETRAZIN SPRAY-ON can be used on all breeds of sheep. VETRAZIN SPRAY-ON is suitable for use on sheep between 4 and 12 weeks after shearing.

A shorter period of protection may occur if VETRAZIN SPRAY-ON is applied incorrectly, if there is excessive fleece soiling or if wool infections such as mycotic dermatitis (fleece rot) are present. Prolonged heavy rain following treatment may also reduce the protection period provided by VETRAZIN SPRAY-ON.

STORAGE

Store in original container, tightly closed, away from direct sunlight, in a cool, dry place, avoiding temperatures below 0°C and above 35°C.

WARRANTY

The manufacturer of this animal remedy extends/grants to the purchaser a warranty that this animal remedy is reasonably fit for the purposes for which its use is recommended, provided that the purchaser uses the remedy only for the purposes for which it is recommended him and strictly in accordance with the directions on this container.

HELPLINE:

Phone 0800 ELANCO (0800 352 626)

Elanco, Vetrazin and the diagonal bar are trademarks owned or licensed by Eli Lilly and Company, its subsidiaries or affiliates.

Registered pursuant to the ACVM Act 1997, No. A005717.

See www.foodsafety.govt.nz for registration conditions.

Registered to Elanco Animal Health (Division of Eli Lilly and Company (NZ) Limited), Level 1, 123 Ormiston Road, Botany Junction, Auckland 2016.

Approved pursuant to the HSNO Act 1996, Approval Code: HSR001844.

See www.epa.govt.nz for approval conditions.

BATCH No.:	
EXPIRY:	

NET CONTENT 25 Litres



9 414644 101732

YL400872NZB

