

**Product name:** Flubenol (A008127)  
**Revision date:** 30 Jul 2020  
**Pack sizes:** 600 g  
**Prepared by:** Paulina Rodriguez  
**Reason:** NZ label, (remove export information)

**Page:** 1 of 7  
**Type of label:** Tube/leaflet  
**Country:** New Zealand



Text above this line is not included in the actual label.

---

**LABEL- FRONT**

**CAUTION  
KEEP OUT OF REACH OF CHILDREN  
FOR ANIMAL TREATMENT ONLY  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

**FLUBENOL™ 5%**

Active Ingredient: 50 mg/g Flubendazole

**Oral wormer for pigs and poultry**

**Net Content: 600g.**

**Elanco Logo**

**Batch No.  
Expiry Date**

**Product name:** Flubenol (A008127)  
**Revision date:** 30 Jul 2020  
**Pack sizes:** 600 g  
**Prepared by:** Paulina Rodriguez  
**Reason:** NZ label, (remove export information)

**Page:** 2 of 7  
**Type of label:** Tube/leaflet  
**Country:** New Zealand

Text above this line is not included in the actual label.

---

***BASE Label***

---

**FLUBENOL™ 5%**

**ORAL WORMER FOR PIGS AND POULTRY**

Flubendazole 50 mg/g

600g

Elanco Australasia Pty Ltd

[www.elanco.co.nz](http://www.elanco.co.nz)

Ph: 0800 426 633.



Flubenol, Elanco and the diagonal bar logo are trademarks of Elanco or its affiliates.

**Product name:** Flubenol (A008127)  
**Revision date:** 30 Jul 2020  
**Pack sizes:** 600 g  
**Prepared by:** Paulina Rodriguez  
**Reason:** NZ label, (remove export information)

**Page:** 3 of 7  
**Type of label:** Tube/leaflet  
**Country:** New Zealand

Text above this line is not included in the actual label.

---

## **MULTI-FOLD LEAFLET**

---

# **FLUBENOL™ 5%**



Active Ingredient: 50 mg/g Flubendazole

## **Oral wormer for pigs and poultry**

**Flubenol 5% is indicated for worm infestations in pigs and poultry.**

**Pigs:** infestations with gastrointestinal worms (*Ascaris suum*, *Hyostrogylus rubidus*, *Oesophagostomum dentatum*, *Trichuris suis*, *Strongyloides ransomi*) and lungworms (*Metastrongylus apri*).

**Poultry:** infestations with gastrointestinal worms (*Ascaridia galli*, *Heterakis gallinarum*, *Capillaria spp.*, *Amidostomum anseris* and *Trichostrongylus tenuis*, *Raillietina sp.*) and gapeworm (*Syngamus trachea*).

## **DOSAGE**

### **PIGS**

**For treatment of piglets and fatteners:** 30g flubendazole per ton of feed (30ppm) for 5 consecutive days in case of a severe infestation with *T. suis*, the animals should be treated for 10 days.

**For treatment of breeding pigs:** 30g flubendazole per ton of feed (30ppm) during 10 consecutive days.

### **Individual and single administration to breeding pigs:**

Flubenol 5% is strewn over the feed in a dose of 5mg per kg of body weight. This corresponds to 1 measuring spoon per 130kg body weight.



**Product name:** Flubenol (A008127)  
**Revision date:** 30 Jul 2020  
**Pack sizes:** 600 g  
**Prepared by:** Paulina Rodriguez  
**Reason:** NZ label, (remove export information)

**Page:** 4 of 7  
**Type of label:** Tube/leaflet  
**Country:** New Zealand

Text above this line is not included in the actual label.

## POULTRY

**Turkeys:** 20g flubendazole per ton of feed (20ppm) for 7 consecutive days.

**Chickens and geese:** 30g flubendazole per ton of feed (30 ppm) for 7 consecutive days.

In case of infestation with tapeworm (*Rallietina sp.*) give 60g flubendazole per ton of feed (60ppm) for 7 consecutive days.

### **Pheasants and partridges:**

60g flubendazole per ton of feed (60ppm) for 7 consecutive days.

## DIRECTIONS FOR USE

Flubenol 5% should be mixed into the feed and the mixture administered.

For a group treatment, make sure that all animals take in enough Flubenol 5%. The medicine should therefore be well mixed into the feed so that a homogenous mixture is obtained. It should not be strewn over pellets or crumbs since this gives a nonhomogeneous mixture.

- 20g flubendazole per ton (20ppm) corresponds to 400g of Flubenol 5% per ton of feed.
- 30g flubendazole per ton (30ppm) corresponds to 600g of Flubenol 5% per ton of feed.
- 60g flubendazole per ton (60ppm) corresponds to 1.2kg of Flubenol 5% per ton of feed.

For small numbers of animals, 1 measuring spoon of Flubenol 5% is mixed into the following amounts of compound feed for the following species: pheasants and partridges: 11kg; chickens and geese: 21.5 kg; turkeys: 32kg.

### **Important:**

Treatment with Flubenol 5% gives optimal results only when strict hygiene of the pig sheds and the poultry houses is practiced.



**Product name:** Flubenol (A008127)  
**Revision date:** 30 Jul 2020  
**Pack sizes:** 600 g  
**Prepared by:** Paulina Rodriguez  
**Reason:** NZ label, (remove export information)

**Page:** 5 of 7  
**Type of label:** Tube/leaflet  
**Country:** New Zealand

Text above this line is not included in the actual label.

## CONTRAINDICATIONS

The use of Flubenol 5% is contraindicated in pigeons and parrots.

## SIDE EFFECTS

In pigs an overdose in the feed of 250 ppm flubendazole or higher (i.e. more than 8 times the normal dose) can bring about short-lasting diarrhea (soft or watery dung), without other clinical signs or effects with reference to the growth of the animals. This loose dung occurs from the second treatment day and is maximal after around 7-12 days.

In poultry even 3 times the highest dose (180 ppm flubendazole) for 7 days had no negative effects on egg production, egg quality, hatching results growth of offspring or in any other respect.

**By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.**

## WITHHOLDING PERIODS:

### PIGS

MEAT: Pigs producing meat or offal for human consumption must not be sold for slaughter either during treatment or within 7 days of the last treatment.

### CHICKENS

MEAT: Chickens producing meat or offal for human consumption must not be sold for slaughter either during treatment or within NIL days of the last treatment.

EGGS: Eggs from treated birds must not be sold for human consumption for NIL days following the last treatment.

### TURKEYS AND GEESE

MEAT: Turkeys and geese producing meat or offal for human consumption must not be sold for slaughter either during treatment or within 1 day of the last treatment.

### GAME BIRDS

MEAT: Game birds producing meat or offal for human consumption must not be sold for slaughter either during treatment or within 4 days of the last treatment



**Product name:** Flubenol (A008127)  
**Revision date:** 30 Jul 2020  
**Pack sizes:** 600 g  
**Prepared by:** Paulina Rodriguez  
**Reason:** NZ label, (remove export information)

**Page:** 6 of 7  
**Type of label:** Tube/leaflet  
**Country:** New Zealand

Text above this line is not included in the actual label.

**It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.**

**Safety Directions:** In connection with potential hypersensitivity reactions and contact dermatitis when mixing, handling and/or administering the product, direct skin contact with the product and inhalation of the product should be avoided. For this purpose, the person administering the product should wear gloves and a dust mask.

**First Aid:** For advice contact the National Poisons Centre **0800 POISONS (0800 764766)** or a doctor immediately.

**WARNING:** May cause damage to organs through prolonged or repeated oral exposure.

Read label before use. Do not breathe dust. Get medical advice or attention if you feel unwell.

**Environmental protection:** Very toxic to aquatic life. DO NOT contaminate wetlands or watercourses with this product or used containers.

**Disposal:** Preferably dispose of the product by use. Otherwise dispose of product and packaging at an approved landfill or other approved facility. Dispose in accordance with local requirements.

For further details see the Safety Data Sheet.

**STORAGE:** Flubenol 5% should be kept at 15-30C (Room temperature).  
Feed prepared with Flubenol 5% can be kept for up to 8 weeks.

**CONDITIONS OF SALE:** The manufacturer of this animal remedy extends/grants to the purchaser a warranty that this animal remedy is reasonably fit for the purposes for which its use is recommended, provided that the purchaser uses the remedy only for the purpose for which it is recommended and strictly in accordance with the directions on this container.

Registered Pursuant to the ACVM Act 1997, No. A006818. See [www.foodsafety.govt.nz/acvm](http://www.foodsafety.govt.nz/acvm) for registration conditions.

**Company Confidential**

---

**Product name:** Flubenol (A008127)  
**Revision date:** 30 Jul 2020  
**Pack sizes:** 600 g  
**Prepared by:** Paulina Rodriguez  
**Reason:** NZ label, (remove export information)

**Page:** 7 of 7  
**Type of label:** Tube/leaflet  
**Country:** New Zealand

Text above this line is not included in the actual label.

---

Registered to  
Elanco Australasia Pty Ltd  
27/88 Shortland St  
Auckland, 1010.



**Phone 0800 426 633**

FLUBENOL, Elanco and the diagonal bar logo are trademarks of Elanco or its affiliates.

Batch No.  
Expiry

**Net Contents: 600g**