

Fly, Maggot and Lice Eliminator for Sheep

Up to 12 weeks Fly Prevention

WARNING KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY



ACTIVE INGREDIENTS: 12.5 g/L Spinosad, 500 g/L cyromazine



10 LITRES

makes 5000 litres of dipwash for Jetting or CR Shower Application

L28320413

PARASITICIDES

YL086829NZX

Fly, Maggot and Lice Eliminator for Sheep



PEEL
HERE
FOR LABEL DIRECTIONS

WARNING Read the label before use
May be harmful if swallowed
Spinosad and cyromazine possibly may cause target organ damage through prolonged or repeated oral exposure
Harmful to aquatic life with long lasting effects
Harmful to the soil environment
Very toxic to terrestrial invertebrates

Action: The active ingredients in Cyrex are spinosad and cyromazine. Spinosad is a member of the "Spinosyn" class of insecticides; a new generation insecticide with a unique mode of action. Cyromazine is an insect growth regulator that kills maggots by stopping their development.

Safety Precautions and Directions

Avoid skin contact. Wash hands and exposed skin before meals and after work. Do not rehandle sheep until dry after treatment.

FIRST AID

If medical advice is needed, have product container or label at hand. Call the National Poisons Centre 0800 POISON (0800 764 766) or a doctor if you feel unwell.

If swallowed do not induce vomiting.

Environmental Protection

Avoid contamination of any water supply or release to the environment of the product or empty container.

DISPOSAL:

Preferably dispose of product by use. Otherwise dispose of product as part of an approved chemical disposal program. Triple rinse empty container and add rinsate to dipwash. Dispose of empty triple-rinsed container as part of an approved plastic recycling program. Otherwise crush and bury empty container at an approved landfill. Do not use container for any other purpose.

EMERGENCY RESPONSE

In case of spillage wear appropriate protective clothing and prevent material from entering waterways. Absorb spills with inert material and place in waste containers. Wash area with water and absorb with further inert material. Dispose of waste safely. For specialist advice in an emergency call 0800 CHEMCALL (0800 243 622).

WITHHOLDING PERIODS

It is an offence for users of this product to cause residues exceeding the relevant MRL in the NZ (Maximum Residues of Agricultural Compounds) Food Standard.

MEAT: Sheep producing meat and offal for human consumption must not be sold for slaughter either during treatment or within 7 days of the last treatment.

MILK: Milk intended for sale for human consumption must be discarded during treatment for not less than 35 days following the last treatment.

Wool Residues: Ensure the time between dipping and subsequent shearing is at least 2 months, except in the case of spot treatment for flystrike.

Cyrex Liquid: New science for high potency, clean wool and operator safety.

Store below 30°C in a dry place in a tightly closed container. Do not freeze.

Do not store in direct sunlight.

Registered to: Elanco Animal Health

(A Division of Eli Lilly and Company (NZ) Ltd.)

Level 1, 123 Ormiston Road, Botany Junction, Auckland, 2016

Registered pursuant to the ACVM Act 1997, No A09917. See www.foodsafety.govt.nz for registration conditions

For advice on your dipping programme, contact your supplier or call the Elanco Helpline: 0800 ELANCO (352 626)

Elanco, Cyrex and the diagonal bar are trademarks owned or licensed by Eli Lilly and Company, its subsidiaries or affiliates.

BATCH NUMBER

EXPIRY



YL086829NZX

L28320413



DIRECTIONS FOR USE:

By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

For the prevention and treatment of blowfly (including *Lucilia cuprina*) strike in all breeds of sheep, including organophosphate resistant strains, for up to 12 weeks.

For the control of lice on coarse wool breeds, including strains resistant to synthetic pyrethroids.

For the treatment of lice on long wool Merino sheep (greater than 3 months wool), including strains resistant to synthetic pyrethroids.

JETTING

Mix 2 litres of Cyrex with 1000 litres of water. Ensure saturation to skin level. Top up at the same rate.

Apply a minimum of 2 litres of dipwash per sheep and an additional 0.5 litres for each month of wool growth up to a maximum of 5 litres per sheep.

CR SHOWER

Mix 2 litres of Cyrex with 1000 litres of water. Ensure saturation to skin level. Top up at the same rate.

No reinforcement is required with the use of a Constant Replenishment (CR) Shower Dip. Do not allow the sump to fall below ¼ of total volume.

FLYSTRIKE DRESSING

Mix 10ml of Cyrex with 5 litres of water. Apply onto the affected area for the spot treatment of flystrike. Remove the wool from around the area to be treated using clippers or shears. Prepare a fresh dilution of flystrike dressing each day.

GENERAL INFORMATION

Fly

A shorter period of protection may occur if Cyrex is applied incorrectly, if there is excessive fleece soiling or if dags present.

Lice

For effective control of lice, thorough wetting is essential, check skin wetting using a wetting pencil. Optimum lice control requires the application of good agricultural practice. This includes attention to stock security following treatment, quarantine treatment of new stock arrivals, thorough and accurate dipping management and utilising the principles of Integrated Pest Management recommended by Wool Pro.

Fly and Lice

It is good agricultural practice not to treat animals when they are wet or if rain is likely. Prolonged heavy rain following treatment may reduce efficacy. Consider retreatment of sheep after heavy rain to ensure continued protection.

MIXING DIRECTIONS:

SHAKE CONTAINER WELL UNTIL PRODUCT IS FULLY MIXED

Mix well, preferably by re-circulating the dip through the pump before use.

Prepare fresh dip, jetting fluid or fly-strike dressing daily. At the end of the day, discard unused product.



WARNING

Read the label before use
May be harmful if swallowed
Spinosad and cyromazine possibly may cause target organ damage through prolonged or repeated oral exposure
Harmful to aquatic life with long lasting effects
Harmful to the soil environment
Very toxic to terrestrial invertebrates

Action: The active ingredients in Cyrex are spinosad and cyromazine. Spinosad is a member of the "Spinosyn" class of insecticides; a new generation insecticide with a unique mode of action. Cyromazine is an insect growth regulator that kills maggots by stopping their development.

Safety Precautions and Directions

Avoid skin contact. Wash hands and exposed skin before meals and after work. Do not rehandle sheep until dry after treatment.

FIRST AID

If medical advice is needed, have product container or label at hand. Call the National Poisons Centre 0800 POISON (0800 764 766) or a doctor if you feel unwell. If swallowed do not induce vomiting.

Environmental Protection

Avoid contamination of any water supply or release to the environment of the product or empty container.

DISPOSAL:

Preferably dispose of product by use. Otherwise dispose of product as part of an approved chemical disposal program. Triple rinse empty container and add rinsate to dipwash. Dispose of empty triple-rinsed container as part of an approved plastic recycling program. Otherwise crush and bury empty container at an approved landfill. Do not use container for any other purpose.

EMERGENCY RESPONSE

In case of spillage wear appropriate protective clothing and prevent material from entering waterways. Absorb spills with inert material and place in waste containers. Wash area with water and absorb with further inert material. Dispose of waste safely. For specialist advice in an emergency call 0800 CHEMCALL (0800 243 622).

WITHHOLDING PERIODS

It is an offence for users of this product to cause residues exceeding the relevant MRL in the NZ (Maximum Residues of Agricultural Compounds) Food Standard.

MEAT: Sheep producing meat and offal for human consumption must not be sold for slaughter either during treatment or within 7 days of the last treatment.

MILK: Milk intended for sale for human consumption must be discarded during treatment for not less than 35 days following the last treatment.

Wool Residues: Ensure the time between dipping and subsequent shearing is at least 2 months, except in the case of spot treatment for flystrike.

Cyrex Liquid: New science for high potency, clean wool and operator safety.

Store below 30°C in a dry place in a tightly closed container. Do not freeze.

Do not store in direct sunlight.

Registered to: Elanco Animal Health

(A Division of Eli Lilly and Company (NZ) Ltd.)

Level 1, 123 Ormiston Road, Botany Junction, Auckland, 2016

Registered pursuant to the ACVM Act 1997, No A09917. See www.foodsafety.govt.nz for registration conditions

For advice on your dipping programme, contact your supplier or call the Elanco Helpline: 0800 ELANCO (352 626)

Elanco, Cyrex and the diagonal bar are trademarks owned or licensed by Eli Lilly and Company, its subsidiaries or affiliates.