

DIRECTIONS FOR USE: FOR INTRAMUSCULAR AND SLOW INTRAVENOUS USE.

DOSAGE:

Cattle: 5 to 10 mg/kg body weight daily.
Pigs: 5 to 10 mg/kg body weight daily.
Sheep and Goats: 10 mg/kg body weight daily.

WITHHOLDING PERIODS:

It is an offence for users of this product to cause residues exceeding the relevant MRL in the NZ (Maximum Residues of Agricultural Compounds) Food Standard.

Meat: Animals producing meat and offal for human consumption must not be sold for slaughter either during treatment or within 21 days of the last treatment.

Milk: Milk intended for sale for human consumption or manufacture for human consumption must be discarded during treatment and for 72 hours following the last treatment.

CAUTION AND SIDE EFFECTS:

Read enclosed cautions and side effects before use.

WARNING: May cause an allergic skin reaction. Causes eye irritation

Safety precautions and directions

Wear protective gloves. Wash hands thoroughly after handling.
Contaminated clothing should not be allowed out of the workplace.
Wash contaminated clothing before reuse.

First Aid: IF IN EYES: Rinse cautiously with water for several minutes.
Remove contact lenses, if present and easy to do. Continue rinsing.
If eye irritation persists, get medical advice.

IF ON SKIN: Wash with plenty of soap and water. If skin irritation or rash occurs, get medical advice. For advice, contact the National Poisons Centre 0800 POISON (0800 764766) or a doctor.

**FOR ANIMAL TREATMENT ONLY
KEEP OUT OF REACH OF CHILDREN
RESTRICTED VETERINARY MEDICINE**

Elanco™ AH0206

Tylan® 200

(Tylosin)
For Injection

250 ml

Each mL contains 200 mg tylosin activity.

**FOR INTRAMUSCULAR USE IN CATTLE, PIGS,
SHEEP AND GOATS.**

FOR SLOW INTRAVENOUS USE IN CATTLE.

Read Insert directions before use.

**Elanco Animal Health
(Division of Eli Lilly & Company (NZ) Limited)
Level 1, 123 Ormiston Road, Botany Junction,
Auckland 2016.**



Registered pursuant to the ACVM Act 1997, No A907
See www.foodsafety.govt.nz for registration conditions.
Approved pursuant to the HSNO Act 1996, No HSR002368
Registered users of the Trademarks Tylan and Elanco.

INDICATIONS:

Success has been reported in the use of **Tylan** for the control of the following disease conditions:

Cattle: Respiratory infections, Foot-rot (necrotic pododermatitis), Calf diphtheria, Metritis, Acute mastitis.

Sheep and Goats: The treatment of the early stages of the peracute and acute forms of contagious agalactia caused by *Mycoplasma agalactiae* and caprine pleuropneumonia caused by *Mycoplasma mycoides var Capri (M. Capri)*.

Pigs: Swine dysentery and enteritis associated with the presence of *Campylobacter coli* and other organisms sensitive to Tylosin, Swine Erysipelas, Pneumonia and Arthritis due to *Mycoplasma*.

Store below 30°C.

Elanco™, Tylan® and the diagonal bar are trademarks of Eli Lilly and Company.

YL1125DENZX (V01-08-2011)

Batch No:

Expiry Date: